



# Deciphering the nitrate signaling pathway leading to a reduction of primary root growth in *Medicago truncatula*

Submitted by Elisabeth Planchet on Thu, 09/27/2018 - 10:27

Titre	Deciphering the nitrate signaling pathway leading to a reduction of primary root growth in <i>Medicago truncatula</i>
Type de publication	Communication
Type	Communication sans actes dans un congrès
Année	2018
Langue	Anglais
Date du colloque	20-21/09/2018
Titre du colloque	Journées Jeunes Chercheurs BAP
Auteur	Zang, Lili [1], Morère-Le Paven, Marie-Christine [2], Clochard, Thibault [3], Beucher, Daniel [4], Aubry, Catherine [5], Limami, Anis M. [6], Montrichard, Françoise [7]
Pays	France
Ville	Paris
Mots-clés	nitrate signal [8], Nox Documents [9], peroxidase [10], primary root [11], Ros [12], SOD [13]
Résumé en anglais	<p>In the model legume <i>Medicago truncatula</i>, nitrate has been shown to inhibit primary root growth through the reduction of root cell elongation. Nitrate, as an essential nutrient, also acts as a signal molecule that is sensed and transduced through a nitrate transporter MtNPF6.8, with RNAi mutants deficient in MtNPF6.8 being insensitive to nitrate [1, 2]. We tested here whether reactive oxygen species (ROS) could be downstream mediators of the nitrate signal since ROS are able to transduce ABA signal in other contexts and also govern the primary root growth. Thus, we analyzed the distribution of ROS (<math>H_2O_2</math>, <math>O_2^{\bullet-}</math>, <math>\bullet OH</math>) and peroxidase activity all along the primary root of seedlings sensitive or insensitive to nitrate using different genotypes of <i>M. truncatula</i>, three wild types and a npf6.8RNAi mutant grown with or without nitrate, to determine whether nitrate modifies ROS and peroxidase patterns. We found that nitrate modified the morphology of the root tip, induced an increase in <math>H_2O_2</math>, and a decrease in <math>O_2^{\bullet-}</math> and <math>\bullet OH</math> in seedlings sensitive to nitrate (R108, A17, and DZA315-16), but not in seedlings insensitive to nitrate (npf6.8RNAi mutant). These results suggest that ROS and peroxidases are downstream mediators in the nitrate signaling pathway. The origin of the change in ROS accumulation in response to nitrate was further investigated following the activity of major enzymes (peroxidase, SOD, Nox) able to interfere with ROS accumulation.</p>
URL de la notice	<a href="http://okina.univ-angers.fr/publications/ua17616">http://okina.univ-angers.fr/publications/ua17616</a> [14]
Lien vers le document en ligne	<a href="https://journees.inra.fr/jjc-bap-2018">https://journees.inra.fr/jjc-bap-2018</a> [15]

---

## Liens

- [1] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=28541>
- [2] <http://okina.univ-angers.fr/mariechristine.lepaven/publications>
- [3] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=11433>
- [4] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=17267>
- [5] <http://okina.univ-angers.fr/c.aubry/publications>
- [6] <http://okina.univ-angers.fr/m.limami/publications>
- [7] <http://okina.univ-angers.fr/f.montrichard/publications>
- [8] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=25442>
- [9] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=25446>
- [10] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=25444>
- [11] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=25443>
- [12] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=1274>
- [13] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=25445>
- [14] <http://okina.univ-angers.fr/publications/ua17616>
- [15] <https://journees.inra.fr/jjc-bap-2018>

Publié sur *Okina* (<http://okina.univ-angers.fr>)